



(19)

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 729 758 A2**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
04.09.1996 Bulletin 1996/36

(51) Int Cl.<sup>6</sup> **A61K 31/505**(21) Application number: **96300931.1**(22) Date of filing: **12.02.1996**

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT  
SE

(72) Inventor: Yuhpyng, Chen L.  
Waterford, Connecticut 06385 (US)


(30) Priority: **02.03.1995 US 397527**

(74) Representative: Hayles, James Richard et al  
Pfizer Limited,  
Patents Department,  
Ramsgate Road  
Sandwich Kent CT13 9NJ (GB)

(71) Applicant: **PFIZER INC.**  
New York, N.Y. 10017 (US)

(54) **Pyrazolopyrimidines and pyrrolopyrimidines for treatment of neuronal and other disorders**

(57) **Pyrazolopyrimidines and pyrrolopyrimidines of the general formula**



wherein A, X, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are defined in the application, and pharmaceutically acceptable salts of such compounds, are useful in the treatment or prevention of certain neuronal and other disorders.

**EP 0 729 758 A2**

6



1

14

20

2

2


2

1

1

## Summary of the invention

This invention relates to a method of treating, preventing or inhibiting a disorder selected from head traumas, spinal cord trauma, ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia), excitotoxic neuronal damage, epilepsy, stroke, stress induced immune dysfunctions, (e.g., porcine stress syndrome, obsessive-compulsive disorder, bovine shipping fever, equine paroxysmal fibrillation, phobias and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs), phobias, muscular spasms, Parkinson's disease, Huntington's disease, urinary incontinence, senile dementia of the Alzheimer's type, multiinfarct dementia, amyotrophic lateral sclerosis, chemical dependencies and addictions (e.g., dependencies on alcohol, cocaine, heroin, benzodiazepines, or other drugs), and hypoglycemia in a mammal, including a human, comprising administering to such mammal an amount of a compound of the formula




wherein

X is nitrogen or -CR<sup>6</sup>;

A is -NR<sup>1</sup>R<sup>2</sup>, -CR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -C(CR<sup>2</sup>R<sup>12</sup>)R<sup>1</sup>, -NHCR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -OCR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -SCR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -NHNR<sup>1</sup>R<sup>2</sup>, -CR<sup>2</sup>R<sup>11</sup>NHR<sup>1</sup>, -CR<sup>2</sup>R<sup>11</sup>OR<sup>1</sup>, -CR<sup>2</sup>R<sup>11</sup>SR<sup>1</sup>, or -C(O)R<sup>2</sup>;

R<sup>1</sup> is hydrogen, or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with from one to two substituents independently selected from the group consisting of hydroxy, fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy,




C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), amino, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>2</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)C(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHC(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -C(=O)(C<sub>1</sub>-C<sub>4</sub> alkyl),

-C(=O)NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -C(=O)N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl),

-SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), and wherein each of the foregoing C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>1</sup> may contain one or two double or triple bonds;

R<sup>2</sup> is C<sub>1</sub>-C<sub>12</sub> alkyl, aryl or -(C<sub>1</sub>-C<sub>10</sub> alkylene)aryl wherein said aryl is phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidyl, imidazolyl, furanyl, benzofuranyl, benzothiazolyl, isothiazolyl, benzisothiazolyl, thiazolyl, isoxazolyl, benzisoxazolyl, benzimidazolyl, triazolyl, pyrazolyl, pyrrolyl, indolyl, pyrrolopyridyl, oxazolyl, or benzoxazolyl; 3- to 8-membered cycloalkyl or -(C<sub>1</sub>-C<sub>6</sub> alkylene) cycloalkyl, wherein one or two of the carbon atoms of any of said cycloalkyl moieties may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>4</sub> alkanoyl, and wherein R<sup>2</sup> may optionally be substituted with from one to three substituents independently selected from chloro, fluoro and C<sub>1</sub>-C<sub>4</sub> alkyl, or by one substituent selected from hydroxy, bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy,



N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>2</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)-C(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHC(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -C O(C<sub>1</sub>-C<sub>4</sub> alkyl).


-C NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -C N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl),

-SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), and -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), and wherein each of the foregoing C<sub>1</sub>-C<sub>12</sub> alkyl and C<sub>1</sub>-C<sub>10</sub> alkylene moieties in the definition may optionally contain one to three double or triple bonds; or


R<sup>1</sup> and R<sup>2</sup>, taken together with the atom to which they are attached, may form a saturated 3- to 8-membered ring which, if it is a 5- to 8-membered ring, may optionally contain one or two double bonds, and wherein one or two of the carbon atoms of said 5- to 8-membered ring may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkanoyl or benzyl;

R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, fluoro, chloro, bromo, iodo, hydroxy, amino, -O(C<sub>1</sub>-C<sub>6</sub> alkyl), -NH(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -S(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), or -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), wherein each of the foregoing C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>3</sup> may contain one double or triple bond and may optionally be substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy, C<sub>1</sub>-C<sub>3</sub> alkoxy, fluoro, chloro and C<sub>1</sub>-C<sub>3</sub> thioalkyl;

R<sup>4</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy, formyl, -NH(C<sub>1</sub>-C<sub>6</sub> alkyl), -N(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SO<sub>n</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), wherein n is 0, 1 or 2, cyano, hydroxy, carboxy, or amido, wherein each of the foregoing C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>4</sup> may optionally be substituted with one substituent selected from hydroxy, trifluoromethyl, amino, carboxy, amido,




-NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl),




C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>3</sub> thioalkyl, fluoro, bromo, chloro, iodo, cyano and nitro;

R<sup>5</sup> is phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidyl, imidazolyl, furanyl, benzofuranyl, benzothiazolyl, isothiazolyl, benzoisothiazolyl, thiazolyl, benzisoxazolyl, benzisoxazolyl, benzimidazolyl, tritoly, pyrazolyl, pyrrolyl, indolyl, pyrrolopyridyl, benzoxazolyl, oxazolyl, pyrrolidinyl, thiazolidinyl, morpholinyl, piperidinyl, piperazinyl, tetrazolyl, or a 3- to 8-membered cycloalkyl or 9- to 12-membered bicycloalkyl ring, wherein one or two of the carbon atoms in said ring may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkanoyl, phenyl or benzyl, and wherein each of the above R<sup>5</sup> groups may optionally be substituted with one or more substituents, preferably with two or three substituents, independently selected from fluoro, chloro, bromo, formyl, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy and trifluoromethyl, or with one substituent selected from hydroxy, iodo, cyano, nitro, amino, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -COO(C<sub>1</sub>-C<sub>4</sub> alkyl), -CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SO<sub>2</sub>NH<sub>2</sub>, -NH<sub>2</sub>SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>6</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), and wherein each of the foregoing C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>5</sup> may optionally be substituted with from one to two substituents independently selected from fluoro, chloro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, amino, methylamino, dimethylamino and acetyl, and wherein each of the foregoing C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>5</sup> may optionally contain one double or triple bond; with the proviso that R<sup>5</sup> is not unsubstituted phenyl;

R<sup>6</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy, formyl, amino, -NH(C<sub>1</sub>-C<sub>6</sub> alkyl), N(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SO<sub>n</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), wherein n is 0, 1 or 2, cyano, carboxy, or amido, and wherein each of the foregoing (C<sub>1</sub>-C<sub>6</sub>)alkyl moieties in the definition of R<sup>6</sup> may be optionally substituted with one substituent selected from hydroxy, trifluoromethyl, amino, carboxy, amido,



-NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl),



C<sub>1</sub>-C<sub>3</sub> thioalkyl, fluoro, bromo, chloro, iodo, cyano and nitro;  
R<sup>11</sup> is hydrogen, hydroxy, fluoro, chloro, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl), cyano, or -CO(C<sub>1</sub>-C<sub>2</sub> alkyl); and  
R<sup>12</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl;

with the proviso that: (1) when X is -CR<sup>6</sup>, A is not straight chain alkyl; (2) when X is CR<sup>6</sup> and R<sup>5</sup> is unsubstituted cycloalkyl and R<sup>3</sup> and R<sup>4</sup> are both hydrogen and R<sup>6</sup> is hydrogen or methyl, then A is not NHR<sup>2</sup> wherein R<sup>2</sup> is benzyl or thienylmethyl, and (3) when X is -CR<sup>6</sup> and R<sup>5</sup> is p-bromophenyl and R<sup>3</sup>, R<sup>4</sup> and R<sup>6</sup> are methyl, then A is not methylamino or hydroxyethylamino;

and when X is nitrogen, with the further proviso that:

(a) A is not straight chain C<sub>1</sub>-C<sub>12</sub> alkyl;

(b) R<sup>5</sup> is not a sugar group;

(c) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and R<sup>5</sup> is chlorophenyl, then A is not -NH-CH(CH<sub>3</sub>)-(CH<sub>2</sub>)<sub>3</sub>-N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>;

(d) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and A is -NR<sup>1</sup> R<sup>2</sup> wherein R<sup>1</sup> is C<sub>3</sub>-C<sub>7</sub> cycloalkyl, and R<sup>2</sup> is C<sub>2</sub>-C<sub>6</sub> alkenyl, phenyl-(C<sub>1</sub>-C<sub>6</sub> alkylene) or hetero-(C<sub>1</sub>-C<sub>6</sub> alkylene) wherein the hetero radical is furyl, thienyl or pyridinyl, and wherein said phenyl may be substituted with one or more substituents independently selected from fluoro, chloro, bromo and iodo, then R<sup>5</sup> is not tetrahydrofuranlyl or tetrahydropyranylyl;

(e) when R<sup>3</sup> is methoxy, methylthio, or methylsulfonyl; R<sup>4</sup> is hydrogen, and R<sup>5</sup> is tetrahydrofuranlyl or tetrahydropyranylyl, then A is not -NH(C<sub>1</sub>-C<sub>2</sub> alkyl), morpholinyl, hydrazino, or -NHC<sub>2</sub>H<sub>4</sub>C<sub>6</sub>H<sub>5</sub>, the phenyl of which may be substituted by one methyl or two methoxy groups;

(f) when R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, chloro, bromo, -SH, or -S-(C<sub>1</sub>-C<sub>4</sub> alkyl), R<sup>4</sup> is hydrogen and R<sup>5</sup> is C<sub>3</sub>-C<sub>6</sub> cycloalkyl, then A is not hydrazino, -NH(C<sub>1</sub>-C<sub>2</sub> alkyl) or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>12</sub> alkyl);

(g) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and A is -NH(CH<sub>2</sub>)<sub>m</sub> COOH wherein m is 1-12, then R<sup>5</sup> is not phenyl substituted by one fluoro, chloro, bromo or iodo group;

(h) when R<sup>3</sup> is hydrogen, hydroxy, methylthio, chloro or -NHbenzyl, R<sup>4</sup> is hydrogen, and R<sup>5</sup> is chlorophenyl or bromophenyl, then A is not -NH(C<sub>1</sub>-C<sub>12</sub> alkyl), -NHallyl, or -N(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>12</sub> alkyl), wherein said C<sub>1</sub>-C<sub>12</sub> alkyl may be substituted by -NC<sub>2</sub>H<sub>5</sub>, or -NHbenzyl which may be substituted by one or two bromo, chloro, fluoro, -NC<sub>2</sub>H<sub>5</sub> phenyl or morpholinopropyl groups;

(i) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and R<sup>5</sup> is nitrophenyl, then A is not -NHR<sup>2</sup> wherein R<sup>2</sup> is phenyl, benzyl or C<sub>1</sub>-C<sub>12</sub> alkyl which may be substituted by one or two hydroxy groups;

(j) when R<sup>3</sup> is chloro or -O(C<sub>1</sub>-C<sub>6</sub> alkyl), R<sup>4</sup> is hydrogen, and A is -NR<sup>1</sup> R<sup>2</sup> wherein R<sup>1</sup> and R<sup>2</sup> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, then R<sup>5</sup> is not chlorophenyl; and

(k) when R<sup>3</sup> is hydrogen, A is benzyl or phenethyl, and R<sup>4</sup> is fluoro, chloro, bromo or iodo, then R<sup>5</sup> is not 5'-deoxy-ribofuranosyl or 5'-amino-5'-deoxyribofuranosyl;

or a pharmaceutically acceptable salt thereof, that is effective in treating or preventing such disorder. The foregoing method is hereinafter also referred to as "Method A".

The term "halo", as used herein, unless otherwise indicated, includes chloro, fluoro, bromo and iodo.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

The term "one or more substituents," as used herein, includes from one to the maximum number of substituents possible based on the number of available bonding sites.

The compounds of formula I have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers and all stereoisomers of compounds of the formula I, and mixtures thereof.

Formula I above includes compounds identical to those depicted but for the fact that one or more hydrogen, nitrogen or carbon atoms are replaced by radioactive isotopes thereof. Such radiolabeled compounds are useful as research and diagnostic tools in metabolism pharmacokinetic studies and in binding assays.

"Ischemic neuronal damage" includes, for example, neuronal damage resulting from CNS surgery, open heart surgery or procedures during which the function of the cardiovascular system is compromised.

"Stress induced immune dysfunctions" include deficiencies in the immune system in both man and animals that are stress induced. For example, such dysfunctions include (but are not limited to) porcine stress syndrome, obsessive-compulsive disorder, bovine shipping fever, equine paroxysmal fibrillation, phobias and dysfunctions induced by confinement stress in livestock animals (e.g., chicken, cows, pigs, sheep, etc.), sheering stress in sheep or human-animal interaction related stress in dogs. (See DeRoeth, L., "Stress and Disease in Domestic Animals", pp. 285 et al. in Annals New York Acc. Sci., 1993; Liptrap, R.M., "Stress and Reproduction in Domestic Animals", pp. 275-282 in Annals New York Acc. Sci., 1993; and Dallaire, A., "Stress and Behavior in Domestic Animals", p. 269-273 in Annals New York Acc. Sci., 1993).

This invention also relates to a method of treating or preventing a disorder selected from head trauma, spinal cord trauma, ischemic neuronal damage (e.g., cerebral ischemia such as cerebral hippocampal ischemia), excitotoxic neuronal damage, epilepsy, stroke, stress induced immune dysfunctions, phobias, muscular spasms, Parkinson's disease, Huntington's disease, urinary incontinence, senile dementia of the Alzheimer's type, multifactorial dementia, amyotrophic lateral sclerosis, chemical dependencies and additions (e.g., dependencies on alcohol, cocaine, heroin, benzodiazepines, or other drugs), and hypoglycemia in a mammal, including a human, comprising administering to such mammal a CRF receptor antagonizing amount of a compound of the formula I, as depicted and defined above, or a pharmacaceutically acceptable salt thereof. The foregoing method is hereinafter also referred to as "Method B".

Preferred embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) R<sup>1</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, -(C<sub>2</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>4</sub> alkyl), or C<sub>2</sub>-C<sub>4</sub> hydroxyalkyl; (b) R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenylethyl, or benzyl substituted with one or two substituents independently selected from chloro, fluoro, methyl, ethyl, methoxy, ethoxy and t-butyl, or by one trifluoromethyl group; (2-thienyl)methyl; (2-thienyl)ethyl; (2-furanyl)methyl; 2-(4-chlorothieryl)methyl; (2-benzofuranyl)methyl; (2-benzothieryl)methyl; (2-thiazolyl)methyl; or (2-benzothiazolyl)methyl; (c) R<sup>3</sup> is hydrogen, methyl, ethyl, methoxy, fluoro or chloro; (d) R<sup>4</sup> is hydrogen, methyl, ethyl, or n-propyl; and (e) R<sup>5</sup> is phenyl substituted with two or three substituents.

Other preferred embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is CR<sup>6</sup> and: (a) R<sup>1</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, -(C<sub>2</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>4</sub> alkyl), or C<sub>2</sub>-C<sub>4</sub> hydroxyalkyl; (b) R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenylethyl, or benzyl substituted with one or two substituents independently selected from chloro, fluoro, methyl, ethyl, methoxy, ethoxy and t-butyl, or by one trifluoromethyl group; (2-thienyl)methyl; (2-thienyl)ethyl; (2-furanyl)methyl; 2-(4-chlorothieryl)methyl; (2-benzofuranyl)methyl; (2-benzothieryl)methyl; (2-thiazolyl)methyl; or (2-benzothiazolyl)methyl; (c) R<sup>3</sup> is hydrogen, methyl, ethyl, methoxy, fluoro or chloro; (d) R<sup>4</sup> is hydrogen, methyl, ethyl, or n-propyl; and (e) R<sup>5</sup> is phenyl substituted with two or three substituents; and (f) R<sup>6</sup> is hydrogen, methyl, ethyl or chloro; with the proviso that R<sup>2</sup> and R<sup>6</sup> are not both hydrogen.

Other preferred embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is CR<sup>6</sup> and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one of hydroxy, fluoro and C<sub>1</sub>-C<sub>2</sub> alkoxy, and may optionally contain one double or triple bond; (b) R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl moiety of said benzyl may optionally be substituted with one fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkoxy; (c) R<sup>3</sup> is methyl, ethyl, fluoro, chloro or methoxy; (d) R<sup>4</sup> and R<sup>5</sup> are independently selected from hydrogen, methyl and ethyl; and (e) R<sup>6</sup> is phenyl substituted with two or three substituents, said substituents being independently selected from fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>4</sub> alkoxy, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one hydroxy group, C<sub>1</sub>-C<sub>4</sub> alkoxy and fluoro and which may optionally contain one double or triple bond, -(C<sub>1</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>2</sub> alkyl), C<sub>1</sub>-C<sub>3</sub> hydroxyalkyl, hydroxy, formyl, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl), or -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl).

Other preferred embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one of hydroxy, fluoro and C<sub>1</sub>-C<sub>2</sub> alkoxy, and may optionally contain one double or triple bond; (b) R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl moiety of said benzyl may optionally be substituted with one fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkoxy; (c) R<sup>3</sup> is methyl, ethyl, fluoro, chloro or methoxy; (d) R<sup>4</sup> and R<sup>5</sup> are independently selected from hydrogen, methyl and ethyl; and (e) R<sup>6</sup> is phenyl substituted with two or three substituents, said substituents being independently selected from fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>4</sub> alkoxy, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one hydroxy group,

C<sub>1</sub>-C<sub>4</sub> alkoxy and fluoro and which may optionally contain one double or triple bond, -(C<sub>1</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>2</sub> alkyl), C<sub>1</sub>-C<sub>3</sub> hydroxyalkyl, hydroxy, formyl, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl), or -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl).

Other embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted by one hydroxy, fluoro or C<sub>1</sub>-C<sub>2</sub> alkoxy group, and which may optionally contain one double or triple bond, and R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, and wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl moiety of said benzyl may optionally be substituted with fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkoxy; or (b) A is -CR<sup>1</sup>R<sup>2</sup> where R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one C<sub>1</sub>-C<sub>6</sub> alkoxy or hydroxy group, R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl moiety of said benzyl may optionally be substituted by one C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, fluoro, chloro or bromo group, and R<sup>1</sup> is hydrogen or fluoro.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is -(C<sub>1</sub>-C<sub>4</sub> alkylene)aryl, wherein said aryl is phenyl, thienyl, benzoturanlyl, furanyl, benzothienyl, thiazolyl, pyridyl or benzothiazolyl.

Other specific embodiments of this invention include methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is a benzyl group that is optionally substituted on the phenyl moiety with one ethyl, t-butyl, methoxy, trifluoromethyl, nitro, fluoro chloro, or methyl group.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is attached through a methylene or ethylene bridge to quinolyl, pyrrolyl, pyrrolidinyl, pyridyl, tetrahydropyranyl, cyclopropyl, piperidinyl, or benzyl-piperidinyl.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or CR<sup>2</sup> and R<sup>1</sup> and R<sup>2</sup> are, independently, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one substituent selected from hydroxy, methoxy, ethoxy, chloro, fluoro, -OC(O)CH<sub>3</sub>, -OC(O)NHCH<sub>3</sub>, and -C(O)NH<sub>2</sub>.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or CR<sup>2</sup> and R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one substituent selected from methoxy and ethoxy.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and A is -NR<sup>1</sup>R<sup>2</sup> or -CHR<sup>1</sup>R<sup>2</sup> in which R<sup>1</sup> and R<sup>2</sup>, together with the N or CH to which they are attached, form a 5- or 6-membered ring in which one of the ring carbon atoms may optionally be replaced by an oxygen or sulfur atom, e.g., pyrrolidinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, thiazolyl, isoxazolyl, thiadiazolyl, oxadiazolyl, pyridyl, pyrazinyl or pyrimidyl.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or CR<sup>2</sup> and A is -NHCHR<sup>1</sup>R<sup>2</sup> or -OCHR<sup>1</sup>R<sup>2</sup> in which -CHR<sup>1</sup>R<sup>2</sup> is a 5- or 6-membered ring in which one of the ring carbon atoms may optionally be replaced by an oxygen or sulfur atom, e.g., tetrahydrofuranlyl, tetrahydrothiuranlyl and cyclopentanylyl.

Other specific embodiments of this invention include Methods A and B, as described above, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound selected from the following:

- 3-((4-methyl-benzyl)-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amino)-propan-1-ol;
- diethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- 2-(butyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amino)-ethanol;
- dibutyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- butyl-ethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- butyl-ethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- butyl-cyclopropylmethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- di-1-propyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- diallyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;
- butyl-ethyl-[6-chloro-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;

butyl-ethyl-[6-methoxy-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 propyl-ethyl-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4,6-d]pyrimidin-4-yl]-amine;  
 4-(1-ethyl-propyl)-6-methyl-3-methylsulfanyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidine;  
 2-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-ylamino]-butan-1-ol;  
 [3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-(1-methylpropyl)amine;  
 4-(1-methoxymethylpropoxy)-3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidine;  
 n-butyl-ethyl-[2,6-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 di-n-propyl-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 ethyl-n-propyl-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 diethyl-2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 n-butyl-ethyl-[2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 2-[N-n-butyl-N-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amino]-ethanol;  
 4-(1-ethyl-propyl)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 n-butyl-ethyl-[2,5-dimethyl-7-(2,4-dimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(1-ethylpropyl)amine;  
 2-(4-bromo-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-(S)-[7-(4-bromo-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 4-(1-ethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 4-(1-methoxymethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 4-(1-ethyl-butyl)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 [7-(4-bromo-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(1-methoxymethyl-propyl)-amine;  
 2-[7-(2-bromo-4,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-[7-(4-ethyl-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-[7-(2-ethyl-4,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol; and  
 2-[7-(2-fluoromethyl-4,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol.

Whenever reference is made to alkyl, this includes both straight and branched chain alkyl.

Whenever reference is made herein to 3-to 8-membered cycloalkyl or 9- to 12-membered bicycloalkyl containing one to three of O, S or N-Z, it is understood that the oxygen and sulfur ring atoms are not adjacent to each other. The three membered cycloalkyl has just one O, S or N-Z. An example of a six-membered cycloalkyl having O and N is morpholinyl.

Whenever R<sup>2</sup> or R<sup>5</sup> is a heterocyclic group, the attachment of the group is through a carbon atom.

Whenever reference is made herein to C<sub>1</sub>-C<sub>4</sub> alkyl or C<sub>1</sub>-C<sub>6</sub> alkyl which "may contain one or two double or triple bonds" in the definitions of R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup>, it is understood that at least two carbons are present in the alkyl for one double or triple bond, and at least four carbons for two double and triple bonds.

Whenever an alkoxy group, e.g., in the definitions of R<sup>1</sup> and R<sup>2</sup>, may have a double or triple bond, it is understood that such double or triple bond is not directly attached to the oxygen.

Formula I above is intended to include all stereoisomers (e.g., all geometric and optical isomers) as well as all racemates of all individual compounds within the depicted genus.

#### Detailed Description of the Invention

Compounds of the formula I wherein X is nitrogen and the pharmaceutically acceptable acid addition of such compounds, as well as methods for preparing such compounds and salts, are referred to in World Patent Application PCT/US 93/11333, which was filed on November 26, 1993.

Compounds of the formula I wherein X is CR<sup>6</sup> and the pharmaceutically acceptable acid addition of such compounds, as well as methods of preparing such compounds and salts, are referred to in World Patent Application PCT/US 93/10715, which was filed on 12 November 1993.

For use in carrying out the methods of this invention, compounds of the formula I and their pharmaceutically acceptable salts may be administered orally, topically or parenterally. They may be administered alone or in combination with pharmaceutically acceptable carriers, in either single or multiple, e.g. up to three, doses. Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solution and various organic solvents. The pharmaceutical compositions formed by combining the novel compounds of formula I and the pharmaceutically acceptable carriers are then readily administered in a variety of dosage forms such as tablets, powders, lozenges, syrups, injectable solutions and the like. These pharmaceutical compositions can, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus, for purposes of oral administration, tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrants such as starch, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone,



sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tabletting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Preferred materials for this include lactose or milk sugar and high molecular weight polyethylene glycols. Oral compositions may also be administered in the form of a gel. When aqueous suspensions or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if desired, emulsifying or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin and combinations thereof.

For parenteral administration, solutions of the novel compound of formula I in sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solution may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.


Additionally, it is possible to administer the compounds employed in the method of the present invention topically when treating inflammatory conditions of the skin and this may be done by way of creams, jellies, gels, pastes and ointments in accordance with standard pharmaceutical practice.

The compounds of formula I and the pharmaceutically acceptable salts thereof will generally be administered from one to three times per day (i.e., from one to three doses per day), with each dose containing from about 0.1 to about 100 mg/kg body weight, although variations will necessarily occur depending upon the weight and condition of the subject being treated, the nature and severity of the disorder for which the subject is being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 1.0 mg to about 50 mg per kg of body weight per individual dose will most desirably be employed. Variations may nevertheless occur depending upon the species of mammal being treated and the individual subject's response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out.

The methods for testing the compounds for formula I for their CRF antagonist activity are according to the procedures of *Endocrinology*, 116, 1653-1659 (1985) and *Peptides*, 10, 179-188 (1985), which determine the binding affinity of a test compound for a CRF receptor. The binding affinities for the compounds of formula I, expressed as  $IC_{50}$  values, generally range from about 0.2 nanomolar to about 10 micromolar.

## Claims

1. Use of a compound of the formula



wherein


X is nitrogen or -CR<sup>6</sup>,

A is -NR<sup>1</sup>R<sup>2</sup>, -CR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -C(=CR<sup>2</sup>R<sup>12</sup>)R<sup>1</sup>, -NHC(R<sup>1</sup>R<sup>2</sup>R<sup>11</sup>), -OCR<sup>1</sup>R<sup>2</sup>R<sup>11</sup>, -SC(R<sup>1</sup>R<sup>2</sup>R<sup>11</sup>), -NHN(R<sup>1</sup>R<sup>2</sup>), -CR<sup>1</sup>R<sup>11</sup>NHR<sup>1</sup>, -CR<sup>2</sup>R<sup>11</sup>OR<sup>1</sup>, -CR<sup>2</sup>R<sup>11</sup>SR<sup>1</sup>, or -C(O)R<sup>2</sup>;

R<sup>1</sup> is hydrogen, or C<sub>1</sub>-C<sub>8</sub> alkyl which may optionally be substituted with from one to two substituents independently selected from the group consisting of hydroxy, fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>8</sub> alkoxy, -O-CO-(C<sub>1</sub>-C<sub>8</sub> alkyl), -O-CO-NH(C<sub>1</sub>-C<sub>8</sub> alkyl), -O-CO-N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), amino, -NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -N(C<sub>1</sub>-C<sub>2</sub> alkyl)(C<sub>1</sub>-C<sub>4</sub> alkyl), -S(C<sub>1</sub>-C<sub>8</sub> alkyl), -N(C<sub>1</sub>-C<sub>4</sub> alkyl)CO(C<sub>1</sub>-C<sub>4</sub> alkyl), -NHCO(C<sub>1</sub>-C<sub>4</sub> alkyl), -COOH, -CO-O(C<sub>1</sub>-C<sub>4</sub> alkyl), -CONH(C<sub>1</sub>-C<sub>4</sub> alkyl), -CON(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SH, -CN, -NO<sub>2</sub>, -SO(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>NH(C<sub>1</sub>-C<sub>4</sub> alkyl), -SO<sub>2</sub>N(C<sub>1</sub>-C<sub>4</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), and wherein each of the foregoing C<sub>1</sub>-C<sub>8</sub> alkyl moieties in the definition of R<sup>1</sup> may contain one or two double or triple bonds;

R<sup>2</sup> is C<sub>1</sub>-C<sub>12</sub> alkyl, aryl or -(C<sub>1</sub>-C<sub>10</sub> alkylene)aryl wherein said aryl is phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidyl, imidazolyl, furanyl, benzofuranyl, benzothiazolyl, isothiazolyl, benzisothiazolyl, thiazolyl, isoxazolyl, benzisoxazolyl, benzimidazolyl, triazolyl, pyrazolyl, pyrrolyl, indolyl, pyrrolidyl,

ridyl, oxazolyl, or benzoxazolyl; 3- to 8-membered cycloalkyl or  $-(C_1-C_6 \text{ alkylene})$  cycloalkyl, wherein one or two of the carbon atoms of any of said cycloalkyl moieties may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen,  $C_1-C_4$  alkyl or  $C_1-C_4$  alkanoyl, and wherein R<sup>2</sup> may optionally be substituted with from one to three substituents independently selected from chloro, fluoro and  $C_1-C_4$  alkyl, or by one substituent selected from hydroxy, bromo, iodo,  $C_1-C_6$  alkoxy,




$-SO_2(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NH(C_1-C_4 \text{ alkyl})$ , and  $-SO_2N(S_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ , and wherein each of the foregoing  $C_1-C_{12}$  alkyl and  $C_1-C_{10}$  alkylene moieties in the definition may optionally contain one to three double or triple bonds; or

R<sup>1</sup> and R<sup>2</sup>, taken together with the atom to which they are attached, may form a saturated 3- to 8-membered ring which, if it is a 5- to 8-membered ring, may optionally contain one or two double bonds, and wherein one or two of the carbon atoms of said 5- to 8-membered ring may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkanoyl or benzyl;

R<sup>3</sup> is hydrogen,  $C_1-C_6$  alkyl, fluoro, chloro, bromo, iodo, hydroxy, amino,  $-O(C_1-C_6 \text{ alkyl})$ ,  $-NH(C_1-C_6 \text{ alkyl})$ ,  $-N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-SH$ ,  $-S(C_1-C_4 \text{ alkyl})$ ,  $-SO(C_1-C_4 \text{ alkyl})$ , or  $-SO_2(C_1-C_4 \text{ alkyl})$ , wherein each of the foregoing  $C_1-C_4$  alkyl and  $C_1-C_6$  alkyl moieties in the definition of R<sup>3</sup> may contain one double or triple bond and may optionally be substituted with from 1 to 3 substituents independently selected from the group consisting of hydroxy,  $C_1-C_3$  alkoxy, fluoro, chloro and  $C_1-C_3$  thioalkyl;


R<sup>4</sup> is hydrogen,  $C_1-C_6$  alkyl, fluoro, chloro, bromo, iodo,  $C_1-C_6$  alkoxy, formyl,  $-NH(C_1-C_6 \text{ alkyl})$ ,  $-N(C_1-C_6 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-SO_2(C_1-C_6 \text{ alkyl})$ , wherein n is 0, or 2, cyano, hydroxy, carboxy, or amido, wherein each of the foregoing  $C_1-C_6$  alkyl moieties in the definition of R<sup>4</sup> may optionally be substituted with one substituent selected from hydroxy, trifluoromethyl, amino, carboxy, amido,



$C_1-C_3$  alkoxy,  $C_1-C_3$  thioalkyl, fluoro, bromo, chloro, iodo, cyano and nitro;

R<sup>5</sup> is phenyl, naphthyl, thienyl, benzothienyl, pyridyl, quinolyl, pyrazinyl, pyrimidyl, imidazolyl, furanyl, benzo-furanyl, benzothiazolyl, isothiazolyl, benzoisothiazolyl, thiazolyl, isoxazolyl, benzisoxazolyl, benzimidazolyl, triazolyl, pyrazolyl, pyrrolyl, indolyl, pyrrolopyridyl, benzoxazolyl, oxazolyl, pyrrolidinyl, thiazolidinyl, morpholinyl, piperidinyl, piperazinyl, tetrazolyl, or a 3- to 8-membered cycloalkyl or 9- to 12-membered bicyclicalkyl ring, wherein one or two of the carbon atoms in said ring may optionally be replaced, independently, by O, S or N-Z wherein Z is hydrogen,  $C_1-C_4$  alkyl,  $C_1-C_4$  alkanoyl, phenyl or benzyl, and wherein each of the above R<sup>5</sup> groups may optionally be substituted with one or more substituents, preferably with two or three substituents, independently selected from fluoro, chloro, bromo, formyl,  $C_1-C_6$  alkyl,  $C_1-C_6$  alkoxy and trifluoromethyl, or with one substituent selected from hydroxy, iodo, cyano, nitro, amino,  $-NH(C_1-C_4 \text{ alkyl})$ ,  $-N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-COO(C_1-C_4 \text{ alkyl})$ ,  $-CO(C_1-C_4 \text{ alkyl})$ ,  $-SO_2NH(C_1-C_4 \text{ alkyl})$ ,  $-SO_2N(C_1-C_4 \text{ alkyl})(C_1-C_2 \text{ alkyl})$ ,  $-SO_2NH_2$ ,  $-NHSO_2(C_1-C_4 \text{ alkyl})$ ,  $-S(C_1-C_6 \text{ alkyl})$ ,  $-SO_2(C_1-C_6 \text{ alkyl})$ , and wherein each of the foregoing  $C_1-C_4$  alkyl and  $C_1-C_6$  alkyl moieties in the definition of R<sup>5</sup> may optionally be substituted with from one to two sub-

stituents independently selected from fluoro, chloro, hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy, amino, methylamino, dimethylamino and acetyl, and wherein each of the foregoing C<sub>1</sub>-C<sub>4</sub> alkyl and C<sub>1</sub>-C<sub>6</sub> alkyl moieties in the definition of R<sup>6</sup> may optionally contain one double or triple bond, with the proviso that R<sup>6</sup> is not unsubstituted phenyl; R<sup>6</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>6</sub> alkoxy, formyl, amino, -NH(C<sub>1</sub>-C<sub>6</sub> alkyl), N(C<sub>1</sub>-C<sub>6</sub> alkyl)(C<sub>1</sub>-C<sub>2</sub> alkyl), -SO<sub>n</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl), wherein n is 0, 1 or 2, cyano, carboxy, or amido, and wherein each of the foregoing (C<sub>1</sub>-C<sub>6</sub>)alkyl moieties in the definition of R<sup>6</sup> may be optionally substituted with one substituent selected from hydroxy, trifluoromethyl, amino, carboxy, amido,



C<sub>1</sub>-C<sub>3</sub> alkoxy, C<sub>1</sub>-C<sub>3</sub> thioalkyl, fluoro, bromo, chloro, iodo, cyano and nitro;

R<sup>11</sup> is hydrogen, hydroxy, fluoro, chloro, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl), cyano, or -CO(C<sub>1</sub>-C<sub>2</sub> alkyl); and

R<sup>12</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl;

with the proviso that: (1) when X is -CR<sup>6</sup>, A is not straight chain alkyl, (2) when X is -CR<sup>6</sup> and R<sup>6</sup> is unsubstituted cycloalkyl and R<sup>3</sup> and R<sup>4</sup> are both hydrogen and R<sup>6</sup> is hydrogen or methyl, then A is not -NHR<sup>2</sup> wherein R<sup>2</sup> is benzyl or thienylmethyl, and (3) when X is -CR<sup>6</sup> and R<sup>6</sup> is p-bromophenyl and R<sup>3</sup>, R<sup>4</sup> and R<sup>6</sup> are methyl, then A is not methylamino or hydroxyethylamino;

and when X is nitrogen, with the further proviso that:

(a) A is not straight chain C<sub>1</sub>-C<sub>12</sub> alkyl;

(b) R<sup>5</sup> is not a sugar group;

(c) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and R<sup>5</sup> is chlorophenyl, then A is not -NH-CH(CH<sub>3</sub>)-(CH<sub>2</sub>)<sub>3</sub>-N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>;

(d) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and A is -NR<sup>1</sup> R<sup>2</sup> wherein R<sup>1</sup> is C<sub>3</sub>-C<sub>7</sub> cycloalkyl, and R<sup>2</sup> is C<sub>2</sub>-C<sub>6</sub> alkenyl, phenyl-(C<sub>1</sub>-C<sub>6</sub> alkylene) or hetero-(C<sub>1</sub>-C<sub>6</sub> alkylene) wherein the hetero radical is furyl, thienyl or pyridinyl, and wherein said phenyl may be substituted by fluoro, chloro, bromo or iodo, then R<sup>5</sup> is not tetrahydrofuranyl or tetrahydropyranyl;

(e) when R<sup>3</sup> is methoxy, methylthio, or methylsulfonyl, R<sup>4</sup> is hydrogen, and R<sup>5</sup> is tetrahydrofuranyl or tetrahydropyranyl, then A is not -NH(C<sub>1</sub>-C<sub>2</sub> alkyl), morpholinyl, hydrazino, or -NHC<sub>2</sub>H<sub>4</sub>C<sub>6</sub>H<sub>5</sub>, the phenyl of which may be substituted by one methyl or two methoxy groups;

(f) when R<sup>3</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, chloro, bromo, -SH, or -S-(C<sub>1</sub>-C<sub>4</sub> alkyl), R<sup>4</sup> is hydrogen and R<sup>5</sup> is C<sub>3</sub>-C<sub>6</sub> cycloalkyl, then A is not hydrazino, -NH(C<sub>1</sub>-C<sub>2</sub> alkyl) or -N(C<sub>1</sub>-C<sub>6</sub> alkyl) (C<sub>1</sub>-C<sub>12</sub> alkyl);

(g) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and A is -NH(CH<sub>2</sub>)<sub>m</sub> COOH wherein m is 1-12, then R<sup>5</sup> is not phenyl substituted by one fluoro, chloro, bromo or iodo group;

(h) when R<sup>3</sup> is hydrogen, hydroxy, methylthio, chloro or -NHbenzyl, R<sup>4</sup> is hydrogen, and R<sup>5</sup> is chlorophenyl or bromophenyl, then A is not -NH(C<sub>1</sub>-C<sub>12</sub> alkyl), -NHallyl, or -N(C<sub>1</sub>-C<sub>6</sub> alkyl) (C<sub>1</sub>-C<sub>12</sub> alkyl), wherein said C<sub>1</sub>-C<sub>12</sub> alkyl may be substituted by -NC<sub>2</sub>H<sub>5</sub>, or -NH benzyl which may be substituted by one or two bromo, chloro, fluoro, -NC<sub>2</sub>H<sub>5</sub> phenyl or morpholinopropyl groups;

(i) when R<sup>3</sup> and R<sup>4</sup> are hydrogen and R<sup>5</sup> is nitrophenyl, then A is not -NHR<sup>2</sup> wherein R<sup>2</sup> is phenyl, benzyl or C<sub>1</sub>-C<sub>12</sub> alkyl which may be substituted by one or two hydroxy groups;

(j) when R<sup>3</sup> is chloro or -O(C<sub>1</sub>-C<sub>6</sub> alkyl), R<sup>4</sup> is hydrogen, and A is -NR<sup>1</sup>R<sup>2</sup> wherein R<sup>1</sup> and R<sup>2</sup> are independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl, then R<sup>5</sup> is not chlorophenyl; and

(k) when R<sup>3</sup> is hydrogen, A is benzyl or phenethyl, and R<sup>4</sup> is fluoro, chloro, bromo or iodo, then R<sup>5</sup> is not 5'-deoxy-ribofuranosyl or 5'-amino-5'-deoxyribofuranosyl;

or a pharmaceutically acceptable salt thereof;

in the manufacture of a medicament for preventing or inhibiting a disorder selected from head traumas, spinal cord trauma, ischemic neuronal damage, excitotoxic neuronal damage, epilepsy, stroke, stress induced immune dysfunctions, phobias, muscular spasms, Parkinson's disease, Huntington's disease, urinary incontinence, senile dementia of the Alzheimer's type, multiinfarct dementia, amyotrophic lateral sclerosis, chemical dependencies and addictions and hypoglycemia.

2. The use as claimed in claim 1, wherein a dose of the medicament contains a quantity of a compound of formula I, or a pharmaceutically acceptable salt thereof, capable of antagonizing the effects of CRF in a mammal.
3. The use according to claim 1 or claim 2, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) R<sup>1</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, -(C<sub>2</sub>-C<sub>4</sub> alkyne)O(C<sub>1</sub>-C<sub>4</sub> alkyl), or C<sub>2</sub>-C<sub>4</sub> hydroxyalkyl; (b) R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenylethyl, or benzyl substituted with one or two substituents independently selected from chloro, fluoro, methyl, ethyl, methoxy, ethoxy and t-butyl, or with one trifluoromethyl group; (2-thienyl)methyl; (2-thienyl)ethyl; (2-furanyl)methyl; 2-(4-chlorothiophenyl)methyl; (2-benzofuranyl)methyl; (2-benzothienyl)methyl; (2-thiazolyl)methyl; or (2-benzothiazolyl)methyl; (c) R<sup>3</sup> is hydrogen, methyl, ethyl, methoxy, fluoro or chloro; (d) R<sup>4</sup> is hydrogen, methyl, ethyl, or n-propyl; and (e) R<sup>5</sup> is phenyl substituted by two or three substituents.
4. The use according to claim 1 or claim 2, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is -CR<sup>6</sup> and: (a) R<sup>1</sup> is C<sub>1</sub>-C<sub>4</sub> alkyl, -(C<sub>2</sub>-C<sub>4</sub> alkyne)O(C<sub>1</sub>-C<sub>4</sub> alkyl), or C<sub>2</sub>-C<sub>4</sub> hydroxyalkyl; (b) R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, benzyl, phenylethyl, or benzyl substituted with one or two substituents independently selected from chloro, fluoro, methyl, ethyl, methoxy, ethoxy and t-butyl, or with one trifluoromethyl group; (2-thienyl)methyl; (2-thienyl)ethyl; (2-furanyl)methyl; 2-(4-chlorothiophenyl)methyl; (2-benzofuranyl)methyl; (2-benzothienyl)methyl; (2-thiazolyl)methyl; or (2-benzothiazolyl)methyl; (c) R<sup>3</sup> is hydrogen, methyl, ethyl, methoxy, fluoro or chloro; (d) R<sup>4</sup> is hydrogen, methyl, ethyl, or n-propyl; (e) R<sup>5</sup> is phenyl substituted by two or three substituents; and (f) R<sup>6</sup> is hydrogen, methyl, ethyl or chloro; with the proviso that R<sup>2</sup> and R<sup>6</sup> are not both hydrogen.
5. The use according to claim 1 or claim 2, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted by one hydroxy, fluoro or C<sub>1</sub>-C<sub>2</sub> alkoxy group, and which may optionally contain one double or triple bond, and R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl moiety of said benzyl may optionally be substituted with one fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>6</sub> alkoxy group; or (b) A is -CR<sup>1</sup>R<sup>2</sup>R<sup>3</sup> wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one C<sub>1</sub>-C<sub>6</sub> alkoxy or hydroxy group, R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl in said benzyl may optionally be substituted by one C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, fluoro, chloro or bromo group, and R<sup>3</sup> is hydrogen or fluoro.
6. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is -(C<sub>1</sub>-C<sub>4</sub> alkyne)aryl wherein said aryl is phenyl, thienyl, benzofuranyl, furanyl, benzothienyl, thiazolyl, pyridyl or benzothiazolyl.
7. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is benzyl optionally substituted on the phenyl moiety with one ethyl, t-butyl, methoxy, trifluoromethyl, nitro, fluoro, chloro, or methyl group.
8. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and R<sup>2</sup> is attached through a methylene or ethylene bridge to quinolyl, pyrrolyl, pyrrolidinyl, pyridyl, tetrahydropyranyl, cyclopropyl, piperidinyl, or benzyl-piperidinyl.
9. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or -CR<sup>6</sup> and R<sup>1</sup> and R<sup>2</sup> are, independently, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one hydroxy, methoxy, ethoxy, chloro, fluoro, -OC(O)CH<sub>3</sub>, -OC(O)NHCH<sub>3</sub>, or -C(O)NH<sub>2</sub> group.
10. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or -CR<sup>6</sup> and R<sup>2</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl substituted with one substituent selected from methoxy and ethoxy.
11. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and A is -NR<sup>1</sup>R<sup>2</sup> or -CHR<sup>1</sup>R<sup>2</sup> in which R<sup>1</sup> and R<sup>2</sup>, together with the N or CH to which they are attached, form a 5- or 6-membered ring in which one of the ring carbons may optionally be replaced by an oxygen or sulfur atom.

12. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen or CR<sup>6</sup> and A is -NHCHR<sup>1</sup>R<sup>2</sup> or -OCHR<sup>1</sup>R<sup>2</sup> in which CHR<sup>1</sup>R<sup>2</sup> is a 5- or 6-membered ring in which one of the ring carbons may optionally be replaced by an oxygen or sulfur atom.

13. The use according to claim 1 or claim 2, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is nitrogen and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted with one hydroxy, fluoro or C<sub>1</sub>-C<sub>2</sub> alkoxy group and may optionally contain one double or triple bond; (b) R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl in said benzyl may optionally be substituted with fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>2</sub> alkoxy; (c) R<sup>3</sup> is methyl, ethyl, fluoro, chloro or methoxy; (d) R<sup>4</sup> and R<sup>5</sup> are independently selected from hydrogen, methyl and ethyl; and (e) R<sup>6</sup> is phenyl substituted by two or three substituents, said substituents being independently selected from fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>4</sub> alkoxy, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy or fluoro group and which may optionally contain one double or triple bond, -(C<sub>1</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>2</sub> alkyl), C<sub>1</sub>-C<sub>3</sub> hydroxyalkyl, hydroxy, formyl, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl) and -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl).

14. The use according to claim 1 or claim 2, wherein the compound of formula I or the pharmaceutically acceptable salt of such compound that is employed is a compound wherein X is CR<sup>6</sup> and: (a) A is -NR<sup>1</sup>R<sup>2</sup>, -NHCHR<sup>1</sup>R<sup>2</sup>, or -OCHR<sup>1</sup>R<sup>2</sup>, wherein R<sup>1</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, which may optionally be substituted with one hydroxy, fluoro or C<sub>1</sub>-C<sub>2</sub> alkoxy group and may optionally contain one double or triple bond; (b) R<sup>2</sup> is benzyl or C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally contain one double or triple bond, wherein said C<sub>1</sub>-C<sub>6</sub> alkyl or the phenyl in said benzyl may optionally be substituted with fluoro, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>1</sub>-C<sub>2</sub> alkoxy; (c) R<sup>3</sup> is methyl, ethyl, fluoro, chloro or methoxy; (d) R<sup>4</sup> and R<sup>5</sup> are independently selected from hydrogen, methyl and ethyl; and (e) R<sup>6</sup> is phenyl substituted by two or three substituents, said substituents being independently selected from fluoro, chloro, bromo, iodo, C<sub>1</sub>-C<sub>4</sub> alkoxy, trifluoromethyl, C<sub>1</sub>-C<sub>6</sub> alkyl which may optionally be substituted with one hydroxy, C<sub>1</sub>-C<sub>4</sub> alkoxy or fluoro group and which may optionally contain one double or triple bond, -(C<sub>1</sub>-C<sub>4</sub> alkylene)O(C<sub>1</sub>-C<sub>2</sub> alkyl), C<sub>1</sub>-C<sub>3</sub> hydroxyalkyl, hydroxy, formyl, -COO(C<sub>1</sub>-C<sub>2</sub> alkyl), and -C(O)(C<sub>1</sub>-C<sub>4</sub> alkyl).

15. The use according to claim 1 or claim 2, wherein the compound of formula I, or the pharmaceutically acceptable salt of such compound that is employed is a compound selected from the following:

3-[(4-methyl-benzyl)-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amino]-propan-1-ol;  
 diethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 2-[butyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]amino]-ethanol;  
 dibutyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 butyl-ethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 butyl-ethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 butyl-cyclopropylmethyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 di-1-propyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 diallyl-[6-methyl-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 butyl-ethyl-[6-chloro-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 butyl-ethyl-[6-methoxy-3-methylsulfanyl-1-(2,4,6-trichlorophenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 propyl-ethyl-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-amine;  
 4-(1-ethyl-propyl)-6-methyl-3-methylsulfanyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidine;  
 2-[3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-butan-1-ol;  
 [3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-yl]-[1-methylpropyl]amine;  
 4-(1-methoxymethylpropoxy)-3,6-dimethyl-1-(2,4,6-trimethylphenyl)-1H-pyrazolo[3,4-d]pyrimidine;  
 n-butyl-ethyl-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 di-n-propyl-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 ethyl-n-propyl-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 diethyl-2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 n-butyl-ethyl-[2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 2-[N-n-butyl-N-[2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amino]-ethanol;  
 4-(1-ethyl-propyl)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;

n-butyl-ethyl-[2,5-dimethyl-7-(2,4-dimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]amine;  
 2,5-dimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(1-ethylpropyl)amine;  
 2-[7-(4-bromo-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-(S)-[7-(4-bromo-2,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 4-(1-ethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 4-(1-methoxymethyl-propoxy)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 4-(1-ethyl-butyl)-2,5,6-trimethyl-7-(2,4,6-trimethylphenyl)-7H-pyrrolo[2,3-d]pyrimidine;  
 [7-(4-bromo-2,6-dimethyl-phenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-(1-methoxymethyl-propyl)-  
 amine;  
 2-[7-(2-bromo-4,6-dimethyl-phenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-[7-(4-ethyl-2,6-dimethyl-phenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol;  
 2-[7-(2-ethyl-4,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol, and  
 2-[7-(2-fluoromethyl-4,6-dimethylphenyl)-2,5-dimethyl-7H-pyrrolo[2,3-d]pyrimidin-4-ylamino]-butan-1-ol.

16. The use according to claim 1 or claim 2, wherein the medicament is for treating or preventing a stress induced immune dysfunction in a livestock animal that is induced by the stress of confinement.



(19)

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 729 758 A3**

(12)

**EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
29.10.1997 Bulletin 1997/44

(51) Int Cl.<sup>6</sup> **A61K 31/505**

(43) Date of publication A2:  
04.09.1996 Bulletin 1996/36

(21) Application number: **96300931.1**(22) Date of filing: **12.02.1996**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE**

(72) Inventor: Yuhpyng, Chen L.  
Waterford, Connecticut 06385 (US)


(30) Priority: **02.03.1995 US 397527**

(74) Representative: Hayles, James Richard et al  
Pfizer Limited,  
Patents Department,  
Ramsgate Road  
Sandwich Kent CT13 9NJ (GB)

(71) Applicant: **PFIZER INC.**  
New York, N.Y. 10017 (US)

(54) **Pyrazolopyrimidines and pyrrolopyrimidines for treatment of neuronal and other disorders**

(57) Pyrazolopyrimidines and pyrrolopyrimidines of the general formula



wherein A, X, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are defined in the application, and pharmaceutically acceptable salts of such compounds, are useful in the treatment or prevention of certain neuronal and other disorders.

**EP 0 729 758 A3**



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 96 30 0931

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION (IN THE AFTER ALIAS) (Int.Cl.)
D,X	WO 94 13677 A (PFIZER ; CHEN YUHPYNG LIANG (US)) 23 June 1994 * claims 1-12 *	1-16	A61K31/505
D,X	WO 94 13676 A (PFIZER ; CHEN YUHPYNG L (US)) 23 June 1994 * claims 1-12 *	1-16	
P,X	WO 95 10506 A (DU PONT MERCK PHARMA) 20 April 1995 * page 1-4 * * table 10 * * claims 1-10 *	1-16	
P,Y	WO 95 33750 A (PFIZER ; CHEN YUHPYNG L (US)) 14 December 1995 * claims 1-22 *	1-16	
Y	US 4 213 977 A (CREANGE JOHN E ET AL) 22 July 1980 * column 2, line 8 - line 21 *	1-15	
D,X	DE 31 45 287 A (TROPONWERKE GMBH & CO KG) 19 May 1983 * page 10, line 19-21; claims 1-10 *	1-15	
D,X	EP 0 496 617 A (GENSIA PHARMA) 29 July 1992 * page 3, line 31 - line 38; claims 1-27 *	1-15	A61K
X	WO 94 13643 A (PFIZER ; FARACI WILLIAM STEPHEN (US); WELCH WILLARD MCKOWAN JR (US)) 23 June 1994 * claims 1-10 *	1-16	
P,X	EP 0 691 128 A (PFIZER) 10 January 1996 * claims 1-6 * compounds (C) VIII, (D) I	1-15	
		-/--	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 19 August 1997	Examiner Seegert, K
CATEGORY OF CITED DOCUMENTS		1: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application I: document cited for other reasons A: technological background C: non-written disclosure P: intermediate document	
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background C: non-written disclosure P: intermediate document		4: member of the same patent family, corresponding document	

EPT: CHEN YUHPYNG LIANG





European Patent Office

**CLAIMS INCURRING FEES**

The present European patent application comprised at the time of filing more than ten claims

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

**LACK OF UNITY OF INVENTION**

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

See sheet -B-

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 96 30 0931

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.8)
D, Y	US 5 063 245 A (ABREU MARY E ET AL) 5 November 1991 * column 5, line 1 - line 20 * ---	1-16	
D, Y	EP 0 005 205 A (TROPONWERKE GMBH & CO KG) 14 November 1979 * page 7, line 26 - page 8, line 5 * * claims 1-10 * ---	1-15	
Y	PHARM. REV., vol. 43, 1991, pages 425-473, XP000614410 OWENS M.J. ET AL: "Physiology and Pharmacology of Corticotropin-releasing factor" * page 447 * "B. Metabolic Responses to Corticotropin-releasing Factor Administration" * page 457 - page 462 * sections IX and X ---	1-16	
Y	BRAIN RES. (NETHERLANDS), vol. 656, no. 2, 1994, pages 405-408, XP002038091 STRIJBOOS P.J.L.M. ET AL: "Corticotropin-releasing factor antagonist inhibits neuronal damage induced by focal cerebral ischaemia or activation of NMDA receptors in the rat brain" * abstract * ---	1-15	
The present search report has been drawn up for all claims			
Place of search <b>MUNICH</b>		Date of completion of the search <b>19 August 1997</b>	Examiner <b>Seegert, K</b>
CATEGORY OF CITED DOCUMENTS		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application I: document cited for other reasons A: technological background O: non-written disclosure P: intermediate document	
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document			



European Patent  
Office

LACK OF UNITY OF INVENTION  
SHEET B

Application Number

ANNEX

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. The subject-matter of the present application relates to the use of compounds of formula (I) for the manufacture of a medicament for the prevention/treatment of numerous disorders as claimed in Claim 1.

2. The common concept linking the different therapeutic applications as claimed resides in the fact that the compounds exhibit corticotropin-releasing factor (CRF) antagonist activity (see e.g. page 1, lines 18 - 26 of the description).

3. However, this common concept is already known from the prior art (see e.g. WO-A-9413677 or WO-A-9413676, also cited on pages 2 - 3 of the description). Consequently, the therapeutic applications as claimed in the present application are no longer linked by a common concept. Thus the requirements of Article 82 EPC are not met.

4. The following different inventions have been identified:

Use of a compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for preventing or inhibiting a disorder selected from

1. head trauma, spinal cord trauma (partly Claims 1 - 15)
2. ischemic neuronal damage, excitotoxic damage (partly Claims 1 - 15)
3. epilepsy (partly Claims 1 - 15)
4. stroke (partly Claims 1 - 15)
5. stress induced immune dysfunctions (partly Claims 1 - 15, Claim 16)
6. phobias (partly Claims 1 - 15)

European Patent  
OfficeLACK OF UNITY OF INVENTION  
SHEET B

Application Number

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

-2-

7. muscular spasms (partly Claims 1 - 15)
8. Parkinson's disease (partly Claims 1 - 15)
9. Huntington's disease (partly Claims 1 - 15)
10. urinary incontinence (partly Claims 1 - 15)
11. senile dementia of the Alzheimer's type (partly Claims 1 - 15)
12. multiinfarct dementia (partly Claims 1 - 15)
13. amyotrophic lateral sclerosis (partly Claims 1 - 15)
14. chemical dependencies and addictions (partly Claims 1 - 15)
15. hypoglycemia (partly Claims 1 - 15)

5. For the time being the search has been restricted to the first subject-matter.

European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 96 30 0931

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THIS APPLICATION (Int.Cl.6)
Y	BRAIN RES. (NETHERLANDS), vol. 545, no. 1-2, 1991, pages 339-342, XP002038092 LYONS M.K. ET AL: "Corticotropin releasing factor antagonist reduces ischemic hippocampal neuronal injury" * abstract *	1-15	
Y	AM. J. PHYSIOL. (UNITED STATES), vol. 255, no. 3 PT 1, 1988, pages 287-292, XP002038093 WIDMAIER E.P. ET AL: "Regulation of corticotropin-releasing factor secretion in vitro by glucose" * abstract *	1-15	
Y	ANN. PHARMACOTHER. (UNITED STATES), vol. 28, no. 6, 1994, pages 779-791, XP002038094 HAINES S.T. ET AL: "Treatment of infantile spasms" * page 781, right-hand column *	1-15	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Y	CLINICAL NEUROPHARMACOLOGY, vol. 9, no. 2, 1986, pages 202-205, XP002038095 KLAWANS H.L. ET AL: "Myoclonus and Parkinsonism" * abstract *	1-15	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 19 August 1997	Examiner Seegert, K
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure I : intermediate documents		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application I : document cited for other reasons @ : member of the same patent family, corresponding document	

EP 0 729 758 A3 (1997.08.19)